

Local asymptotics for some q -hypergeometric polynomials

Juan F. Mañas Mañas
Universidad de Almería
jmm939@ual.es

Juan J. Moreno-Balcázar
Universidad de Almería
balcazar@ual.es

The basic q -hypergeometric function ${}_r\phi_s$ is defined by the series

$${}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} ; q, z \right) = \sum_{k=0}^{\infty} \frac{(a_1; q)_k \cdots (a_r; q)_k}{(b_1; q)_k \cdots (b_s; q)_k} \left((-1)^k q^{\binom{k}{2}} \right)^{1+s-r} \frac{z^k}{(q; q)_k}, \quad (1)$$

where $0 < q < 1$ and $(a_j; q)_k$ and $(b_j; q)_k$ denote the q -analogues of the Pochhammer symbol.

When one of the parameters a_j in (1) is equal to q^{-n} the basic q -hypergeometric function is a polynomial of degree at most n in the variable z . Our objective is to obtain a type of local asymptotics, known as Mehler–Heine asymptotics, for q -hypergeometric polynomials when $r = s$.

Concretely, by scaling adequately these polynomials we intend to get a limit relation between them and a q -analogue of the Bessel function of the first kind. Originally, this type of local asymptotics was introduced for Legendre orthogonal polynomials (OP) by the German mathematicians H. E. Heine and G. F. Mehler in the 19th century. Later, it was extended to the families of classical OP (Jacobi, Laguerre, Hermite), and more recently, these formulae were obtained for other families as discrete OP, generalized Freud OP, multiple OP or Sobolev OP, among others.

These formulae have a nice consequence about the scaled zeros of the polynomials, i.e. using the well-known Hurwitz’s theorem we can establish a limit relation between these scaled zeros and the ones of a Bessel function of the first kind. In this way, we are looking for a similar result in the context of the q -analysis and we will illustrate the results with numerical examples.