

8ECM, Portorož, June 20-26, 2021

Minisymposium “Variational and evolutionary models involving local/nonlocal interactions”

MINIMAL PLANAR N -PARTITIONS FOR LARGE N

Giovanni Alberti (Università di Pisa)

joint work with

Marco Caroccia (Politecnico di Milano)


Giacomo Del Nin (University of Warwick)

Summary

- ▶ Minimal partitions and Hales's Honeycomb Theorem
- ▶ Uniform energy distribution for minimal partitions
- ▶ **Towards a description of the structure of minimal partitions:
emergence of grains**

Partitions

Let Ω be a two-dimensional domain with finite area.



An N -partition of Ω is a collection $\mathcal{E} = \{E_1, \dots, E_N\}$ of closed sets in Ω (called *cells* of the partition)

- ▶ with pairwise disjoint interiors and union Ω ,
- ▶ with equal area $|E_i| = |\Omega|/N$,
- ▶ and sufficiently regular boundaries.


The *perimeter* of a partition \mathcal{E} is

$$\begin{aligned}\text{Per}(\mathcal{E}) &:= \text{length}(\partial E_1 \cup \cdots \cup \partial E_n) \\ &= \frac{1}{2} \sum_{i=1}^N \text{length}(\partial E_i) + \frac{1}{2} \text{length}(\partial \Omega) \\ &= \frac{1}{2} \sum_{i=1}^N \text{Per}(E_i) + \frac{1}{2} \text{Per}(\Omega)\end{aligned}$$

- ▶ Ω admits a minimal N -partition for every integer N ;
- ▶ the local structure of minimal N -partitions is simple;
- ▶ computing minimal N -partitions is challenging.

Hales's Honeycomb Theorem (T.C. Hales, 2001)

Let Ω be a flat torus which admits a regular hexagonal N -partition \mathcal{E}_{hex} .



$$N=48$$

Then \mathcal{E}_{hex} is the unique minimal N -partition of Ω .

- ▶ Not all flat tori admit a regular hexagonal partition.
- ▶ No counterpart in higher dimension!

Key tool: Hales's isoperimetric inequality

Simplified version (polygons only):

- ▶ let E be an n -polygon with area 1,
- ▶ let R_n be the regular n -polygon with area 1,
- ▶ let $H = R_6$ be the regular hexagon with area 1.

Then:

$$\text{Per}(E) \geq \text{Per}(R_n) \geq \text{Per}(H) - c(n - 6)$$

because $n \mapsto \text{Per}(R_n)$ is a convex function (!)

We can do better:

$$\text{Per}(E) \geq \text{Per}(H) - c(n - 6) + \delta(\text{dist}(E, H))^2$$

where $\text{dist}(E, H)$ is *for example* the Hausdorff distance of E from the closest regular hexagon H .

Idea of proof of Hales's theorem


Let \mathcal{E} be an N -partition of Ω with E_i an n_i -polygon:

$$\begin{aligned}\text{Per}(\mathcal{E}) &= \frac{1}{2} \sum_i \text{Per}(E_i) \\ &\geq N \cdot \frac{1}{2} \text{Per}(H) - \frac{c}{2} \sum_i (n_i - 6) + \frac{\delta}{2} \sum_i (\text{dist}(E_i, H))^2 \\ &= N \cdot \frac{1}{2} \text{Per}(H) + \frac{\delta}{2} \sum_i (\text{dist}(E_i, H))^2 \\ &\geq N \cdot \frac{1}{2} \text{Per}(H) = \text{Per}(\mathcal{E}_{\text{hex}}).\end{aligned}$$

We set $\sigma := \frac{1}{2} \text{Per}(H) = \sqrt[4]{12}$.

Minimal N -partitions of planar domains

We let Ω be an arbitrary planar domain with finite area,



and consider a minimal N -partition \mathcal{E} of Ω with very large N .

- ▶ We expect that most cells are close to regular hexagons
→ local hexagonal patterns.
- ▶ We expect some “disturbance” close to the boundary of Ω .
Does such disturbance decay away from the boundary?
- ▶ Is the orientation of the local hexagonal pattern “constant”?
If not, is it “piecewise constant”? → emergence of “grains”

Uniform energy distribution

- ▶ We can prove a uniform distribution of energy in the spirit of [A. + Choksi + Otto, 2009].
- ▶ Purpose: show that “most” cells are close to be regular hexagons in a quantified way.
- ▶ It is conveniente to replace N with the length parameter

$$\varepsilon := \sqrt{|\Omega|/N};$$


ε^2 is the area of the cells of the partition, now called ε -partition.

Theorem (Energy distribution of the hexagonal partition)

Let \mathcal{E}_{hex} be the regular hexagonal partition of the plane with cells of area 1. Then the “energy density” of \mathcal{E}_{hex} is $\sigma := \sqrt[4]{12}$.

More precisely, for every disc $B = B(x, r)$ with $r \gg 1$:

$$\text{Per}_B(\mathcal{E}_{\text{hex}}) = \sigma \text{area}(B) + O(r^{2/3})$$



- ▶ Statement similar to Gauss's Circle Theorem.
- ▶ Proof by Fourier transform.

By scaling: if $\mathcal{E}_{\text{hex}}^\varepsilon$ is the regular hexagonal ε -partition of the plane, then for every disc $B = B(x, r)$ with $r \gg \varepsilon$

$$\text{Per}_{B}(\mathcal{E}_{\text{hex}}^\varepsilon) = \frac{\sigma}{\varepsilon} \text{area}(B) + O(\varepsilon^{1/3} r^{2/3})$$

Theorem (Uniform distribution of energy)

Let \mathcal{E}_ε be minimal ε -partitions of Ω . Let $B_\varepsilon = B(x_\varepsilon, r_\varepsilon)$ be discs in Ω with $r_\varepsilon \gg \varepsilon$ and $\text{dist}(B_\varepsilon, \partial\Omega) \gg \varepsilon$. Then

$$\text{Per}_{B_\varepsilon}(\mathcal{E}_\varepsilon) = \frac{\sigma}{\varepsilon} |B_\varepsilon| + O(r_\varepsilon).$$

- ▶ Proof of lower bound based Hales inequality.
- ▶ Proof of upper bound based on “cut and paste” technique.
- ▶ A precise statement depends on the variant of the problem considered.

Towards a precise description of minimal ε -partitions

- ▶ Recall the questions: *In the limit $\varepsilon \rightarrow 0$, is the orientation of the local hexagonal pattern constant?
Is it piecewise constant?*
- ▶ Consider the “excess energy”:

$$F_\varepsilon(\mathcal{E}) := \varepsilon \operatorname{Per}(\mathcal{E}) - \sigma|\Omega|;$$


ideally, we would like to write the Γ -limit of F_ε as $\varepsilon \rightarrow 0$.

What should be the variable of such Γ -limit?

The “limit” of the orientation of the local hexagonal patterns.

- ▶ We did not write the Γ -limit, but we identified and partially addressed two key questions (“cell problems”).

Excess energy due to change of orientation




- ▶ Consider a square of side-length $L \gg 1$;
- ▶ consider all 1-partitions \mathcal{E} which are prescribed in the grey zone, as in the picture;
- ▶ $\theta :=$ angle between imposed orientations.

Define

$$\Phi(\theta) := \liminf_{L \rightarrow +\infty} \frac{1}{L} \left\{ \inf_{\mathcal{E}} F_1(\mathcal{E}) \right\}.$$

- ▶ Explicit construction gives $\Phi(\theta) = O(\theta |\log \theta|)$;
- ▶ Is $\Phi(\theta) > 0$? Presumably yes, partial proof.
- ▶ Is $\Phi(\theta)$ superlinear in $\theta = 0$? Presumably yes, no proof.

Excess energy due to boundary



- ▶ Consider a square of side-length $L \gg 1$ and the rectangle R as in the picture;
- ▶ consider all 1-partitions of R which are prescribed in the grey zone, as in the picture;
- ▶ $\theta :=$ angle between the imposed orientation and the vertical direction.

Define

$$\Phi_b(\theta) := \liminf_{L \rightarrow +\infty} \frac{1}{L} \left\{ \inf_{\mathcal{E}} F_1(\mathcal{E}) \right\}.$$

- ▶ Hales isoperimetric inequality gives $C \leq \Phi_b \leq C'$.
- ▶ Does $\Phi_b(\theta)$ depends on θ ? Presumably yes, no proofs.

Conclusions

- ▶ If $\Phi_b(\theta)$ does NOT depend on θ , then minimal ε -partitions of Ω have constant orientation (in the limit $\varepsilon \rightarrow 0$).
- ▶ If $\Phi_b(\theta)$ depends on θ , and Φ is strictly positive then minimal ε -partitions of Ω may not have constant orientation.
- ▶ If in addition $\Phi(\theta)$ is super-linear at 0 then the orientation of minimal ε -partitions is “piecewise constant”.
→ emergence of “grains”.

Thanks for the attention!